

原著

<英文>

Akashi A, Kanamori A, Nakamura M, Fujihara M, Yamada Y, Negi A. Comparative assessment for the ability of Cirrus, RTVue, and 3D-OCT to diagnose glaucoma. *Invest Ophthalmol Vis Sci* 54:4478-84.2013.

Akashi A, Kanamori A, Nakamura M, Fujihara M, Yamada Y, Negi A. The ability of macular parameters and circum papillary retinal nerve fiber layer by three SD-OCT instruments to diagnose highly myopic glaucoma. *Invest Ophthalmol Vis Sci* 54:6025-32.2013

Kanamori A, Naka M, Akashi A, Fujihara M, Yamada Y, Nakamura M. Cluster analyses of grid-pattern display in macular parameters using optical coherence tomography for glaucoma diagnosis. *Invest Ophthalmol Vis Sci* 54:6401-8.2013

Kanamori A, Nakamura M, Tomioka M, Kawaka Y, Yamada Y, Negi A. Structure-function relationship among three types of spectral-domain optical coherent tomography instruments in measuring parapapillary retinal nerve fibre layer thickness. *Acta Ophthalmol* 91:e196-202.2013

Kanamori A, Nakamura M, Yamada Y, Negi A. Spectral-domain optical coherence tomography detects optic atrophy due to optic tract syndrome. *Graefes Arch Clin Exp Ophthalmol* 251:591-5.2013

Miki A, Kanamori A, Negi A, Naka M, Nakamura M. Loss of aquaporin 9 expression adversely affects the survival of retinal ganglion cells. *Am J Pathol* 182:1727-39.2013.

Nakano S, Kanamori A, Nakamura M, Mizukawa K, Negi A. Paraneoplastic optic neuropathy associated with cerebellar choroid meningioma. *Eye (Lond)* 27:1220-1.2013

Nakano S, Kanamori A, Nakamura M, Mizukawa K, Negi A. Paraneoplastic optic neuropathy associated with cerebellar choroid meningioma. *Eye (Lond)* 27:1220-1.2013

Sakamoto M, Kanamori A, Fujihara M, Yamada Y, Nakamura M, Negi A. Assessment of iCareONE rebound tonometer for self-measuring intraocular pressure. *Acta Ophthalmol* 2013.

Tanigawa M, Tsukahara Y, Yamanaka H. A case of acute posterior multifocal placoid pigment epitheliopathy demonstrating vogt-koyanagi-harada disease-like optical coherence tomography findings in the acute stage. *Case Rep Ophthalmol* 4(3):172-9.2013.

Kanamori A, Naka M, Akashi A, Fujihara M, Yamada Y, Nakamura M. Cluster analyses of grid-pattern display in macular parameters using optical coherence tomography for glaucoma diagnosis. *Invest Ophthalmol Vis Sci* 27: 54(9):6401-8. 2013

Akashi A, Kanamori A, Nakamura M, Fujihara M, Yamada Y, Negi A. The ability of macular parameters and circum papillary retinal nerve fiber layer by three SD-OCT instruments to diagnose highly myopic glaucoma. *Invest Ophthalmol Vis Sci* 54(9):6025-32. 2013

Akashi A, Kanamori A, Nakamura M, Fujihara M, Yamada Y, Negi A. Comparative assessment for the ability of Cirrus, RTVue, and 3D-OCT to diagnose glaucoma. *Invest Ophthalmol Vis Sci* 10:54(7):4478-84. 2013

Kanamori A, Nakamura M, Tomioka M, Kawaka Y, Yamada Y, Negi A, Structure-function relationship among three types of spectral-domain optical coherent tomography instruments in measuring parapapillary retinal nerve fibre layer thickness. *Acta Ophthalmol* 91(3):e196-202. 2013

Kanamori A, Nakamura M, Yamada Y, Negi A. Spectral-domain optical coherence tomography detects optic atrophy due to optic tract syndrome. *Graefes Arch Clin Exp Ophthalmol* 251: 591-595.2013

Otsuka H, Kawano H, Sonoda S, Nakamura M, Sakamoto T: Particle-induced endophthalmitis: Possible mechanisms of sterile endophthalmitis after intravitreal triamcinolone. *Invest Ophthalmol Vis Sci* 54: 1758-1766.2013

Miki A, Kanamori A, Negi A, Naka M, Nakamura M. The loss of aquaporin 9 expression adversely affects the survival of retinal ganglion cells. *Am J Pathol* 182: 1727-1739.2013

Sakamoto M, Kanamori A, Fujihara M, Yamada Y, Nakamura M, Negi A. Assessment of IcareOne rebound tonometer for self-measuring intraocular pressure. *Acta Ophthalmol* 2013, Epub ahead of print

Kanamori A, Nakamura M, Tomioka M, Kawaka Y, Yamada Y, Negi A. Structure-function relationship among three types of spectral-domain optical coherent tomography instruments in measuring parapapillary retinal nerve fiber layer thickness. *Acta Ophthalmol* 91: e196-202.2013

Akashi A, Kanamori A, Nakamura M, Fujihara M, Yamada Y, Negi A. Comparative assessment for the ability of Cirrus, RTVue, and 3D-OCT to diagnose glaucoma. *Invest Ophthalmol Vis Sci* 54: 4478-4484.2013.

Kato T, Meguro A, Nomura E, Uemoto R, Nomura N, Ota M, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida K, Inatani M, Tanihira H, Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Chin S, Inoko H, Mizuki N. Association study of genetic variants on chromosome 7q31 with susceptibility to normal tension glaucoma in a Japanese population. *Eye* 27: 979-983.2013.

Kanamori A, Akashi A, Nakamura M, Fujihara M, Yamada Y, Negi A. The ability of macular parameters and circumpapillary retinal nerve fiber layer by three SD-OCT instruments to diagnose highly myopic glaucoma. *Invest Ophthalmol Vis Sci* 54: (9): 6025-6032.2013.

Kanamori A, Naka M, Akashi A, Fujihara M, Yamada Y, Nakamura M. Cluster analyses of grid-pattern display in macular parameters using optical coherence tomography for glaucoma diagnosis. *Invest Ophthalmol Vis Sci* 54 (9): 6401-6408.2013.

Nakano S, Kanamori A, Nakamura M, Mizukawa K, Negi A. Paraneoplastic optic neuropathy associated with cerebellar choroid meningioma. *Eye* 27 (10): 1220-1221.2013.

<和文>

渡部真樹子、坂井 譲：サルコイドーシスによる下垂体鞍上部肉芽腫の1例 臨床眼科 67巻6号:867-873,2013

<緑内障・神経眼科>

根木 昭、中村 誠、金森 章泰、楠原あづさ、中西 賴子、辰巳 康子、藤岡 美幸、田口真衣子、福田 昌秀、田渕久美子、三木 明子、溝上 淳二、長井 隆行、藤原 雅史、山田 裕子：視神経疾患の新しい展開(総説) 日本眼科学会雑誌 117巻3号:187-211,2013

<網膜・硝子体>

下山 剛、横山 知子、野村 耕治：当院における未熟児網膜症および他の新生児合併症の経過 日本眼科学会雑誌 117巻8号:616-620,2013

宇津 永遠、野村 耕治、大西 健、沖本 聰志：当院における未熟児網膜症診療の発症ならびに治療状況 眼科臨床紀要 6(4):306-310,2013.

<斜視・弱視>

沖本 聰志、野村 耕治：調節性内斜視に対する累進多焦点コンタクトレンズの有用性 眼科臨床紀要 6(3):203-207,2013