

Publication list of Makoto Nakamura as of July 14, 2014 (Only written in English)

I.Original Articles

1. Nakamura M, Ara F, Yamada M, Hotta Y, Hayakawa M, Fujiki K, Kanai A, Sakai J, Inoue M, Yamamoto M, Fujiwara Y, Umoto A, Miyazaki S, Shimo-oku M, Furuyama J, Nakajima A, Imachi J. High frequency of mitochondrial ND4 gene mutation in Japanese pedigrees with Leber hereditary optic neuropathy. *Jpn J Ophthalmol* 1992, 36: 56-61. (IF=0.924, CI=24)
2. Nakamura M, Fujiwara Y, Yamamoto M. Homoplasmic and exclusive ND4 gene mutation in Japanese pedigrees with Leber's disease. *Invest Ophthalmol Vis Sci* 1993, 34: 488-495. (IF=3.597, CI=12)
3. Nakamura M, Fujiwara Y, Yamamoto M. The two locus control of Leber hereditary optic neuropathy and a high penetrance in Japanese pedigrees. *Hum Genet* 1993, 91: 339-341. (IF=5.069, CI=29)
4. Fujiwara Y, Nakamura M, Yokoo S. A new anticancer platinum compound, (-)-(R)-2-aminomethylpyrrolidine (1,1-cyclobutanedicarboxylato) platinum (II): DNA interstrand crosslinking, repair and lethal effects in normal human, Fanconi's anaemia and xeroderma pigmentosum cells. *Br J Cancer* 1993, 67:1285-1292. (IF=5.042, CI=10)
5. Nakamura M. Genetic analysis of Japanese pedigrees with Leber's hereditary optic neuropathy. *Kobe Journal of Medical Sciences* 1993, 39:171-182. (IF=0, CI=0)
6. Nakamura M, Yamamoto M. DNA interstrand crosslinking agents and human ocular fibroblasts: Differential sensitivity to Mitomycin-C and cis-Diaminedichloroplatinum (II). *Exp Eye Res* 1994, 59:53-62. (IF=3.259, CI=7)
7. Hotta Y, Fujiki Y, Hayakawa M, Nakajima A, Kanai A, Mashima Y, Hiida Y, Shinoda K, Yamada K, Oguchi Y, Ishida M, Yanashima K, Wakakura M, Ishikawa S, Nakamura M, Sakai J, Yamamoto M, Hayashi T, Mitani I, Miyazaki S, Shimo-oku M, Imachi J, Kuniyoshi N, Nagataki S, Isashiki Y, Ohba N. Clinical features of Japanese Leber's hereditary optic neuropathy with 11778 mutation of mitochondrial DNA. *Jpn J Ophthalmol* 1995, 39: 96-108. (IF= 0.924, CI=33)
8. Nakamura M, Sekiya Y, Yamamoto M. Preservation of photic blink reflex in Leber's hereditary optic neuropathy. *Invest Ophthalmol Vis Sci* 1996, 37: 2736-2743. (IF=3.597, CI=6)
9. Nakamura M, Sekiya Y, Yamamoto M. Evaluation of bilateral undefined optic neuropathy based on mitochondrial genetics. *Neuro-ophthalmology* 1996, 16: 91-97. (IF=0.256, CI=2)
10. Yamamoto M, Dogru M, Nakamura M, Shirabe H, Tsukahara Y, Sekiya Y. Visual function following congenital cataract surgery. *Jpn J Ophthalmol* 1998, 42: 411-416. (IF=0.924, CI=13)
11. Dogru M, Inoue M, Nakamura M, Yamamoto M. Modifying factors related to asymmetric diabetic retinopathy. *Eye* 1998, 12: 929-933. (IF=1.851, CI=14)

12. Shirabe H, Suda K, Mori Y, Dogru M, Nakamura M, Sekiya Y, Yamamoto M. Visual function following unilateral congenital cataract surgery. American Orthoptic Journal 1998, 48: 97-103. (IF=0, CI=0)
13. Yamada K, Oguchi Y, Hotta Y, Nakamura M, Isashiki Y, Mashima Y. Multicenter study on the frequency of three primary mutations of mitochondrial DNA in Japanese pedigrees with Leber's hereditary optic neuropathy: Comparison with American and British counterparts. Neuro-ophthalmology 1999, 22: 187-193. (IF=0.256, CI=9)
14. Dogru M, Nakamura M, Inoue M, Yamamoto M. Long-term visual outcome in proliferative diabetic retinopathy patients after panretinal photocoagulation. Jpn J Ophthalmol 1999, 43: 217-224. (IF=0.924, CI=8)
15. Maeda H, Tanaka Y, Nakamura M, Yamamoto M. Blue-on-Yellow perimetry using an Armaly glaucoma screening program. Ophthalmologica 1999, 213: 71-75. (IF=1.424, CI=7)
16. Maeda H, Nakamura M, Yamamoto M. Morphometric features of laminar pores in lamina cribrosa observed by scanning laser ophthalmoscopy. Jpn J Ophthalmol 1999, 43: 415-421. (IF=0.924, CI=8)
17. Dogru M, Shirabe H, Nakamura M, Sekiya Y, Matsuo M, Taoka K, Nomura K, Yamamoto M. Development of grating acuity in infants with retinopathy of prematurity. Acta Ophthalmol Scand 1999, 77:72-75. (IF=1.848, CI=4)
18. Nakamura M, Hara R, Kimura R, Ohgiya N, Yamamoto N, Morino I, Yamamoto M. Optic perineuritis not associated with syphilitic infection. Neuro-ophthalmology 1999, 21: 135-145. (IF=0.256, CI=2)
19. Maeda H, Eno A, Nakamura M, Negi A. Safe management of a late-onset bleb leak with a needling technique. Eye 2000, 14: 802-804. (IF=1.851, CI=6)
20. Maeda H, Nakamura M, Negi A. New perimetric threshold test algorithm with dynamic strategy and tendency oriented perimetry (TOP) in glaucomatous eyes. Eye 2000, 14: 747-751. (IF=1.851, CI=7)
21. Nakamura M, Yamamoto M. Variable pattern of visual recovery of Leber's hereditary optic neuropathy. Br J Ophthalmol 2000, 84: 534-535. (IF=2.902, CI=16)
22. Dogru M, Shirabe H, Nakamura M, Taoka K, Nomura K, Yamamoto M. Effect of retinopathy of prematurity on resolution acuity development in 1-to 3-year-old children. J Pediatric Ophthalmol Strabismus 2001, 38: 144-148. (IF=0.626, CI=0)
23. Maeda H, Nakamura M, Negi A. Selective reduction of the S-cone component of the electroretinogram in Posner-Schlossman syndrome. Eye 2001, 15:163-167. (IF=1.851, CI=1)
24. Barber AJ, Nakamura M (**equal contributor**), Wolpert EB, Reiter CEN, Seigel GM, Antonetti DA, Gardner TW. Insulin rescues retinal neurons from apoptosis by a phosphatidylinositol

- 3-kinase/akt-mediated mechanism that reduces the activation of caspase-3. *J Biol Chem* 2001, 276: 32814-32821. (IF=4.773, CI=154)
25. Nakamura M, Barber AJ, Antonetti DA, LaNoue KF, Robinson KA, Buse MG, Gardner TW. Excessive hexosamines block the neuroprotective effect of insulin and induce apoptosis in retinal neurons. *J Biol Chem* 2001, 276: 43748-43755. (IF=4.773, CI=73)
26. Kanamori A, Escano MT, Eno A, Nakamura M, Maeda H, Seya R, Ishibashi K, Negi A. Evaluation of the effect of aging on retinal nerve fiber layer thickness measured by optic coherence tomography. *Ophthalmologica* 2003, 217: 273-278. (IF=1.424, CI=55)
27. Kanamori A, Nakamura M, Escano MFT, Seya R, Maeda H, Negi A. Evaluation of the glaucomatous damage on retinal nerve fiber layer thickness measured by optical coherence tomography. *Am J Ophthalmol* 2003, 135:513-520. (IF=4.223, CI=146)
28. Kanamori A, Nakamura M (corresponding author), Mukuno H, Maeda H, Negi A. Diabetes has an additive effect on neural apoptosis in rat retina with chronically elevated intraocular pressure. *Curr Eye Res* 2004, 28: 47-54. (IF=1.28, CI=35)
29. Kanamori A, Matsui N, Tatsumi Y, Nagai A, Nakanishi Y, Nakamura M, Negi A. Optical coherence tomography detects characteristic retinal nerve fiber layer thickness corresponding to band atrophy of the optic discs. *Ophthalmology* 2004, 111: 2278-2283. (IF=5.454, CI=38)
30. Kanamori A, Nakamura M (corresponding author), Nakanishi Y, Nagai A, Mukuno H, Yamada Y, Negi A. Akt is activated via insulin / IGF-1 receptor in rat retina with episcleral vein cauterization. *Brain Res* 2004, 1022: 195-204. (IF=2.728, CI=19)
31. Mukuno H, Nakamura M (corresponding author), Nagai A, Kanamori A, Negi A. Unoprostone isopropyl rescues retinal progenitor cells from apoptosis in vitro. *Curr Eye Res* 2004, 29: 457-464. (IF=1.28, CI=13)
32. Kanamori A, Nagai-Kusuhara A, Escano MF, Maeda H, Nakamura M, Negi A. Comparison of confocal scanning laser ophthalmoscopy, scanning laser polarimetry and optical coherence tomography to discriminate ocular hypertension and glaucoma at an early stage. *Graefes Arch Clin Exp Ophthalmol* 2005, 26: 1-11. (IF=2.17, CI=46)
33. Kanamori A, Nakamura M (corresponding author), Nakanishi Y, Yamada Y, Negi A. Long-term glial reactivity in rat retinas ipsilateral and contralateral to experimental glaucoma. *Exp Eye Res* 2005, 81: 48-56. (IF=3.259, CI=26)
34. Reiter CE, Wu X, Sandirasegarane L, Nakamura M, Gilbert KA, Singh RS, Fort PE, Antonetti DA, Gardner TW. Diabetes reduces basal retinal insulin receptor signaling: reversal with systemic and local insulin. *Diabetes* 2006, 55: 1148-1156. (IF=8.286, CI=48)
35. Nakanishi Y, Nakamura M (corresponding author), Tatsumi Y, Nagai-Kusuhara A, Negi A. Quantification of retinal nerve fiber thickness reduction associated with a relative afferent pupillary defect. *Graefes Arch Clin Exp Ophthalmol* 2006, 244:1480-1484. (IF=2.17 CI=4)

36. Kusuvara S, Nakamura M, Nagai-Kusuvara A, Nakanishi Y, Kanamori A, Negi A. Macular thickness reduction in eyes with unilateral optic atrophy detected with optical coherence tomography. *Eye* 2006; 20: 882-887. (IF=1.851, CI=6)
37. Nakanishi Y, Nakamura M (**corresponding author**), Mukuno H, Kanamori A, Seigel GM, Negi A. Latanoprost rescues retinal neuro-glial cells from apoptosis by inhibiting caspase-3, which is mediated by p44/p42 mitogen-activated protein kinase. *Exp Eye Res* 2006; 83:1108-1117. (IF=3.259, CI=19)
38. Nakamura M, Darhad U, Tatsumi Y, Fujioka M, Kusuvara A, Maeda H, Negi A. Agreement of rebound tonometer in measuring intraocular pressure with three types of applanation tonometers. *Am J Ophthalmol* 2006; 142: 332-334. (IF=4.223, CI=32)
39. Nagai-Kusuvara A, Nakamura M (**corresponding author**), Mukuno H, Kanamori A, Negi A, Seigel GM. cAMP-responsive element binding protein mediates a cGMP/protein kinase G-dependent anti-apoptotic signal induced by nitric oxide in retinal neuro-glial progenitor cells. *Exp Eye Res.* 2007; 84: 152-162. (IF=3.259, CI=13)
40. Fujioka M, Nakamura M (**corresponding author**), Tatsumi Y, Kusuvara A, Maeda H, Negi A. Comparison of Pentacam Scheimpflug camera with ultrasound pachymetry and noncontact specular microscopy in measuring central corneal thickness. *Curr Eye Res* 2007; 32: 89-94. (IF=1.28, CI=21)
41. Tatsumi Y, Nakamura M (**corresponding author**), Fujioka M, Nakanishi Y, Kusuvara A, Maeda H, Negi A. Quantification of retinal nerve fiber layer thickness reduction associated with a relative afferent pupillary defect in asymmetric glaucoma. *Br J Ophthalmol* 2007; 91: 633-637. (IF=2.902, CI=7)
42. Nagai-Kusuvara A, Nakamura M (**corresponding author**), Kanamori A, Nakanishi Y, Kusuvara S, Negi A. Evaluation of optic nerve head configuration in various types of optic neuropathy with Heidelberg Retina Tomograph. *Eye*. 2008; 22: 1154-1160. (IF=1.851, CI=7)
43. Nagai-Kusuvara A, Nakamura M (**corresponding author**), Fujioka M, Tatsumi Y, Negi A. Association of retinal nerve fiber layer thickness measured by confocal scanning laser ophthalmoscopy and optical coherence tomography with disc size and axial length. *Br J Ophthalmol* 2008; 92: 186-190. (IF=2.902, CI=26)
44. Nagai-Kusuvara A, Nakamura M (**corresponding author**), Fujioka M, Negi A. Long-term results of amniotic membrane transplantation-assisted bleb revision for leaking blebs. *Graefes Arch Clin Exp Ophthalmol.* 2008; 246: 567-571. (IF=2.17, CI=6)
45. Shibuya E, Meguro A, Ota M, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida T, Inatani M, Tanihara H, Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Ohno S, Inoko H, Mizuki

- N. Association of Toll-like receptor 4 gene polymorphisms with normal tension glaucoma. Invest Ophthalmol Vis Sci. 2008, 49: 4453-4457. (IF=3.597, CI=25)
46. Nagai-Kusuhara A, Nakamura M (corresponding author), Tatsumi Y, Nakanishi Y, Negi A. Disagreement between Heidelberg Retina Tomograph and optical coherence tomography in assessing optic nerve head configuration of eyes with band atrophy and normal eyes. Br J Ophthalmol. 2008, 92: 1382-1386. (IF=2.902, CI=8)
47. Tatsumi Y, Kanamori A, Nagai-Kusuhara A, Nakanishi Y, Agarwal N, Negi A, Nakamura M (corresponding author). Nipradilol protects rat retinal ganglion cells from apoptosis induced by serum deprivation in vitro and by diabetes in vivo. Curr Eye Res. 2008, 33: 683-692. (IF=1.28, CI=2)
48. Akiyama M, Yatsu K, Ota M, Katsuyama Y, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Kumagai N, Nishida T, Inatani M, Tanihara H, Ohno S, Inoko H, Mizuki N. Microsatellite analysis of the GLC1B locus on chromosome 2 points to NCK2 as a new candidate gene for normal tension glaucoma. Br J Ophthalmol. 2008, 92: 1293-1296. (IF=2.902, CI=4)
49. Darhad U, Nakamura M (corresponding author), Fujioka M, Tatsumi Y, Nagai-Kusuhara A, Maeda H, Negi A. Intraocular pressure lowering effect of once daily versus once weekly latanoprost instillation in the same normal individuals. Kobe J Med Sci. 2008, 53:297-304. (IF=0, CI=0)
50. Kanamori A, Naka M, Nagai-Kusuhara A, Yamada Y, Nakamura M, Negi A. Regional relationship between retinal nerve fiber layer thickness and corresponding visual field sensitivity in glaucomatous eyes. Arch Ophthalmol. 2008, 126: 1500-1506. (IF=3.711, CI=14)
51. Murakami Y, Ikeda Y, Yonemitsu Y, Onimaru M, Nakagawa K, Kohno R, Miyazaki M, Hisatomi T, Nakamura M, Yabe T, Hasegawa M, Ishibashi T, Sueishi K. Inhibition of nuclear translocation of apoptosis-inducing factor is an essential mechanism of the neuroprotective activity of pigment epithelium-derived factor in a rat model of retinal degeneration. Am J Pathol 2008, 173: 1326-1338. (IF=4.89, CI=30)
52. Nakamura M, Kanamori A, Nagai-Kusuhara A, Kusuhara S, Yamada Y, Negi A. Serous macular detachment due to diabetic papillopathy detected using optical coherence tomography. Arch Ophthalmol 2009, 127: 105-107. (IF=3.711, CI=0)
53. Nagai-Kusuhara A, Nakamura M (corresponding author), Kanamori A, Negi A. Association of optic disc configuration and clustered visual field sensitivity in glaucomatous eyes with hemifield visual field defects. J Glaucoma 2009, 18: 62-68. (IF=1.776, CI=2)
54. Kanamori A, Naka M, Fukuda M, Nakamura M, Negi A. Latanoprost protects rat retinal ganglion cells from apoptosis in vitro and in vivo. Exp Eye Res. 2009, 88: 535-541. (IF=3.259, CI=9)

55. Arimura N, Ki-I Y, Hashiguchi T, Kawahara K, Biswas KK, Nakamura M, Sonoda Y, Yamakiri K, Okubo A, Sakamoto T, Maruyama I. Intraocular expression and release of high-mobility group box 1 protein in retinal detachment. *Lab Invest* 2009, 89: 278-289. (IF=3.641, CI=17)
56. Nakamura J, Meguro A, Ota M, Nomura E, Nishide T, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida T, Inatani M, Tanihara H, Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Ohno S, Inoko H, Mizuki N. Association of toll-like receptor 2 gene polymorphisms with normal tension glaucoma. *Mol Vis* 2009, 15: 2905-2910. (IF=2.205, CI=2)
57. Kanamori A, Naka M, Fukuda M, Nakamura M, Negi A. Tafluprost protects rat retinal ganglion cells from apoptosis in vitro and in vivo. *Graefes Arch Clin Exp Ophthalmol.* 2009, 247: 1353-1360. (IF=2.17, CI=6)
58. Nakamura K, Ota M, Meguro A, Nomura N, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida T, Inatani M, Tanihara H, Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Ohno S, Inoko H, Mizuki N. Association of microsatellite polymorphisms of the GPDS1 locus with normal tension glaucoma in the Japanese population. *Clin Ophthalmol* 2009, 3: 307-312. (IF=0, CI=0)
59. Kamino M, Meguro A, Nomura N, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida T, Inatani M, Tanihara H, Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Ohno S, Inoko H, Mizuki N. Investigation of the association between the GLC3A locus and normal tension glaucoma in Japanese patients with microsatellite analysis. *Clin Ophthalmol* 2009, 3: 183-188. (IF=0, CI=0)
60. Naka M, Kanamori A, Tatsumi Y, Fujioka M, Nagai-Kusuhara A, Nakamura M, Negi A. Comparison of mean deviation with AGIS and CIGTS scores in association with structural parameters in glaucomatous eyes. *J Glaucoma* 2009, 18: 379-384. (IF=1.776, CI=1)
61. Suzuki M, Meguro A, Ota M, Nomura E, Kato T, Nomura N, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida T, Inatani M, Tanihara H, Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi. Genotyping HLA-DRB1 and HLA-DQB1 alleles in Japanese patients with normal tension glaucoma. *Mol Vis* 2010, 16: 1874-1879. (IF=2.205, CI=0)
62. The writing committee for the normal tension glaucoma genetic study group of Japan Glaucoma Society. Genome-wide association study of normal tension glaucoma: Common variants in SRBD1 and ELOVL5 contribute to disease susceptibility. *Ophthalmology* 2010, 117: 1331-1338. (IF=5.454, CI=20)
63. Murakami K, Meguro A, Ota M, Shiota T, Nomura N, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida T, Inatani M, Tanihara H,

- Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Ohno S, Inoko H, Mizuki N. Analysis of microsatellite polymorphisms within the GLC1F locus in Japanese patients with normal tension glaucoma. Mol Vis 2010, 16: 462-466. (IF=2.205, CI=0)
64. Fukuda M, Nakanishi Y, Fuse M, Yokoi N, Hamada Y, Fukagawa M, Negi A, Nakamura M (corresponding author). Altered expression of aquaporins 1 and 4 coincides with neurodegenerative events in retinas of spontaneously diabetic Torii rats. Exp Eye Res 2010, 90: 17-25. (IF=3.259, CI=5)
65. Naka M, Kanamori A, Negi A, Nakamura M (corresponding author). Reduced expression of aquaporin-9 in rat optic nerve head and retina following elevated intraocular pressure. Invest Ophthalmol Vis Sci 2010, 51: 4618-4626. (IF=3.597, CI=7)
66. Otsuka H, Arimura N, Sonoda S, Nakamura M, Hashiguchi T, Maruyama I, Nakao S, Hafezi-Moghadam A, Sakamoto T. Stromal cell-derived factor 1 is essential for photoreceptor cell protection in retinal detachment. Am J Pathol 2010, 177: 2268-2277. (IF=4.89, CI=7)
67. Ishikawa K, Nagai T, Yamada Y, Negi A, Nakamura M (corresponding author). Optimal conditions for multifocal VEP recording for normal Japanese population established by receiver operating characteristic analysis. Doc Ophthalmol 2011, 122: 29-37. (IF=2.107, CI=2)
68. Fukuda M, Naka M, Mizokami J, Negi A, Nakamura M (corresponding author). Diabetes induces expression of aquaporin-0 in the retinal nerve fibers of spontaneously diabetic Torii rats. Exp Eye Res 2011, 91: 195-201 (IF=3.259, CI=1).
69. Yamamoto T, Kuwayama Y, The Collaborative Bleb-related Infection Incidence and Treatment Study Group. Interim clinical outcomes in the collaborative bleb-related infection incidence and treatment study. Ophthalmology 2011, 118: 453-458. (IF=5.454, CI=5)
70. Yasumura R, Meguro A, Ota M, Nomura E, Uemoto R, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida T, Inatani M, Tanihara H, Aihara M Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Ohno S, Inoko H, Mizuki N. Investigation of the association between SLC1A3 gene polymorphisms and normal tension glaucoma. Mol Vis 2011, 17: 792-796. (IF=2.205, CI=1)
71. Mizokami J, Yamada Y, Negi A, Nakamura M (corresponding author). Postural changes in intraocular pressure are associated with asymmetrical retinal nerve fiber thinning in treated patients with primary open-angle glaucoma. Graefes Arch Clin Exp Ophthalmol, 2011, 249:879-885. (IF=2.17, CI=3)
72. Fort PE, Losiewicz MK, Reiter CEN, Singh RSJ, Nakamura M, Abcouwer SF, Barber AJ, Gardner TW. Differential roles of hyperglycemia and hypoinsulinemia in diabetes induced retinal cell death: Evidence for retinal insulin resistance. PloS One 2011, 6: e26498. (IF=4.092, CI=1)

73. Nakamura M, Ishikawa K, Nagai T, Negi A. Receiver-operating characteristic analysis of multifocal VEPs to diagnose and quantify glaucomatous functional damage. Doc Ophthalmol 2011, 123: 93-108. (IF=2.107, CI=0)
74. Mizokami J, Kanamori A, Negi A, Nakamura M (**corresponding author**). A preliminary study of reduced expression of aquaporin-9 in the optic nerve of primate and human eyes with glaucoma. Curr Eye Res 2011, 36: 1064-1067. (IF=1.28, CI=1)
75. Nakamura M, Naka M, Tatsumi Y, Nagai-Kusuhara A, Kanamori A, Yamada Y, Negi A. Filtering bleb structure associated with long-term intraocular pressure control after amniotic membrane-assisted trabeculectomy. Curr Eye Res 2012, 37: 239-250. (IF=1.28, CI=0)
76. Kanamori A, Nakamura M, Tabuchi K, Yamada Y, Negi A. Effects of ocular rotation on parapapillary retinal nerve fiber layer thickness analysis measured with spectral-domain optical coherence tomography. Jpn J Ophthalmol. 2012, 56:354-361. (IF=0.924, CI=1)
77. Kanamori A, Nakamura M, Tomioka M, Kawaka Y, Yamada Y, Negi A. Agreement among three types of spectral-domain optical coherent tomography instruments in measuring parapapillary retinal nerve fibre layer thickness. Br J Ophthalmol 2012, 96: 832-837. (IF=2.902, CI=1)
78. Kanamori A, Nakamura M, Yamada Y, Negi A. Longitudinal study of retinal nerve fiber layer thickness and ganglion cell complex in traumatic optic neuropathy. Arch Ophthalmol. 2012, 130:1067-1069. (IF=3.711, CI=0)
79. Nakamura M, Ishikawa-Tabuchi K, Kanamori A, Yamada Y, Negi A. Better performance of RTVue than Cirrus spectral-domain optical coherence tomography in detecting band atrophy of the optic nerve. Graefes Arch Clin Exp Ophthalmol. 2012, 250: 1499-1507. (IF=2.17, CI=0)
80. Kanamori A, Nakamura M, Yamada Y, Negi A. Spectral-domain optical coherence tomography detects optic atrophy due to optic tract syndrome. Graefes Arch Clin Exp Ophthalmol 2013, 251: 591-595. (IF=2.17, CI=0)
81. Otsuka H, Kawano H, Sonoda S, Nakamura M, Sakamoto T: Particle-induced endophthalmitis: Possible mechanisms of sterile endophthalmitis after intravitreal triamcinolone. Invest Ophthalmol Vis Sci 2013; 54: 1758-1766. (IF=3.597, CI=3)
82. Miki A, Kanamori A, Negi A, Naka M, Nakamura M (**corresponding author**). The loss of aquaporin 9 expression adversely affects the survival of retinal ganglion cells. Am J Pathol 2013, 182: 1727-1739 (IF=4.89, CI=0)
83. Sakamoto M, Kanamori A, Fujihara M, Yamada Y, Nakamura M, Negi A. Assessment of IcareOne rebound tonometer for self-measuring intraocular pressure. Acta Ophthalmol 2013, Epub ahead of print (IF=2.629, CI=0)
84. Kanamori A, Nakamura M, Tomioka M, Kawaka Y, Yamada Y, Negi A. Structure-function relationship among three types of spectral-domain optical coherent tomography instruments in

- measuring parapapillary retinal nerve fiber layer thickness. *Acta Ophthalmol* 2013, 91: e196-202 (IF=2.629, CI=0)
85. Akashi A, Kanamori A, Nakamura M, Fujihara M, Yamada Y, Negi A. Comparative assessment for the ability of Cirrus, RTVue, and 3D-OCT to diagnose glaucoma. *Invest Ophthalmol Vis Sci* 2013; 54: 4478-4484. (IF=3.597, CI=0).
86. Kato T, Meguro A, Nomura E, Uemoto R, Nomura N, Ota M, Kashiwagi K, Mabuchi F, Iijima H, Kawase K, Yamamoto T, Nakamura M, Negi A, Sagara T, Nishida K, Inatani M, Tanihira H, Aihara M, Araie M, Fukuchi T, Abe H, Higashide T, Sugiyama K, Kanamoto T, Kiuchi Y, Iwase A, Chin S, Inoko H, Mizuki N. Association study of genetic variants on chromosome 7q31 with susceptibility to normal tension glaucoma in a Japanese population. *Eye* 2013; 27: 979-983. (IF=1.851, CI=0).
87. Akashi A, Kanamori A, Nakamura M, Fujihara M, Yamada Y, Negi A. The ability of macular parameters and circumpapillary retinal nerve fiber layer by three SD-OCT instruments to diagnose highly myopic glaucoma. *Invest Ophthalmol Vis Sci* 2013, 54(9):6025-32. (IF=3.597, CI=0).
88. Kanamori A, Naka M, Akashi A, Fujihara M, Yamada Y, Nakamura M. Cluster analyses of grid-pattern display in macular parameters using optical coherence tomography for glaucoma diagnosis. *Invest Ophthalmol Vis Sci*. 2013, 54(9):6401-8.
89. Matsumoto Y, Kanamori A, Nakamura M, Takahashi T, Nakashima I, Negi A. Sera from patients with seropositive neuromyelitis optica spectral disorders caused the degeneration of rodent optic nerve. *Exp Eye Res*. 2014, 119:61-9.
90. Yamamoto T, Sawada A, Mayama C, Araie M, Ohkubo S, Sugiyama K, Kuwayama Y; Collaborative Bleb-Related Infection Incidence and Treatment Study Group. The 5-year incidence of bleb-related infection and its risk factors after filtering surgeries with adjunctive mitomycin C: collaborative bleb-related infectionincidence and treatment study 2. *Ophthalmology*. 2014, 121(5):1001-6.
91. Matsumoto Y, Fujihara M, Kanamori A, Yamada Y, Nakamura M (**corresponding author**). Effect of axial length reduction after trabeculectomy on the development of hypotony maculopathy. *Jpn J Ophthalmol*. 2014, 58(3):267-75.
92. Matsumoto Y, Kanamori A, Nakamura M, Negi A. Rat chronic glaucoma model induced by intracameral injection of microbeads suspended in sodium sulfate-sodium hyaluronate. *Jpn J Ophthalmol*. 2014 May, (3):290-7.
93. Kawano H, Ito T, Yamada S, Hashiguchi T, Maruyama I, Hisatomi T, Nakamura M, Sakamoto T. Toxic effects of extracellular histones and their neutralization by vitreous in retinal detachment. *Lab Invest*. 2014, 94(5):569-85.

94. Nakamura M, Kato K, Kamata S, Ishikawa K, Nagai T. Effect of refractive errors on multifocal VEP responses and standard automated perimetry tests in a single population. Doc Ophthalmol. 2014, 128(3):179-89.
95. Akashi A, Kanamori A, Ueda K, Matsumoto Y, Yamada Y, Nakamura M. The detection of macular analysis by SD-OCT for optic chiasmal compression neuropathy and nasotemporal overlap. Invest Ophthalmol Vis Sci. 2014, Epub ahead of print.

II. Case Reports

1. Nakamura M, Tanigawa M, Yamamoto M. A case of Leber's hereditary optic neuropathy with a mitochondrial DNA mutation at nucleotide position 3460. Jpn J Ophthalmol 1994, 38: 267-271. (IF=0.924, CI=6)
2. Nakamachi Y, Nakamura M, Fujii S, Yamamoto M, Okubo K. Oguchi disease with sectoral retinitis pigmentosa harboring adenine deletion at position 1147 in the arrestin gene. Am J Ophthalmol 1998, 125: 249-251. (IF=4.223, CI=22)
3. Nakamura M, Tanabe M, Yamada Y, Azumi A. Zoster sine herpete with bilateral ocular involvement. Am J Ophthalmol 2000, 129:809-810. (IF=4.223, CI=8)
4. Nakamura M, Kanamori A, Seya R, Maeda H, Negi A. A case of occult macular dystrophy accompanying normal-tension glaucoma. Am J Ophthalmol 2003, 135:715-717. (IF=4.223, CI=4)
5. Kanamori A, Nakamura M, Matsui N, Tomimori H, Tanase M, Seya R, Maeda H, Negi A. Goniosynechialysis with lens aspiration and posterior chamber intraocular lens implantation for glaucoma in spherophakia. J Cataract Refract Surg 2004, 30: 513-516. (IF=2.2264, CI=4)
6. Nagai A, Kanamori A, Kusuvara S, Nakamura M, Negi A. Unilateral manifestation of Leber's hereditary optic neuropathy after blunt ocular trauma. Jpn J Ophthalmol 2005, 49: 65-67. (IF=0.924, CI=1)
7. Nakamura M, Kanamori A, Kusuvara S, Ishibashi K, Negi A. Alternate total ophthalmoplegia and optic neuropathy associated with ulcerative colitis. Eye 2005, 19: 235-237. (IF=1.851, CI=0)
8. Tatsumi Y, Kanamori A, Kusuvara A, Nakanishi Y, Kusuvara S, Nakamura M (**corresponding author**). Retinal nerve fiber layer thickness in optic tract syndrome. Jpn J Ophthalmol 2005, 49: 294-296. (IF=0.924, CI=7)
9. Kakutani Y, Nakamura M (**corresponding author**), Nagai-Kusuvara A, Kanamori A, Negi A. Marked cup reversal presumably associated with scleral biomechanics in a case of adult glaucoma. Arch Ophthalmol 2010, 128: 139-141. (IF=3.711, CI=3)

10. Nakano S, Kanamori A, Nakamura M, Mizukawa K, Negi A. Paraneoplastic optic neuropathy associated with cerebellar choroid meningioma. *Eye* 2013, Epub ahead pf print (IF=1.851, CI=0)

III. Review

1. Fujiwara Y, Nakamura A, Yokoo S, Nakamura M. Senescence of normal human and progeroid cells as a potential model of aging. In, New horizons in aging science. Eds., Orimo H, Fukuchi Y, Kuramoto K, Iriki M. The University of Tokyo Press, Tokyo, 1992, 87-88.
2. Gardner TW, Antonetti DA, Barber AJ, LaNoue KF, Nakamura M. New insights into the pathophysiology of diabetic retinopathy: potential cell-specific therapeutic targets. *Diabet Technol & Ther* 2000, 2: 601-608. (IF=1.931, CI=25).
3. Barber AJ, Nakamura M, Penn State Retina Research Group. Apoptosis and neurodegeneration in diabetes: Lessons from the retina. In, Diabetic renal-retinal syndrome. Eds., Friedman EA, L'Esperance FA Jr. Kluwer Academic Publishers, Netherlands, 2002, 35-45.
4. Nakamura M, Kanamori A, Negi A. Diabetes mellitus as a risk factor for glaucomatous optic neuropathy. *Ophthalmologica* 2005, 219: 1-10 (IF=1.424, CI=29).